

CHAPTER 23
ENVIRONMENTAL
CHEMISTRY

After completing this lesson, you will be able to:

- Recognize various chemical reactions occurring in the atmosphere. (Understanding)
- Recognize that the release of COX SOX NOX VOCs is associated with the combustion of hydrocarbon-based fuels. (Applying)
- Outline problem associated with release of pollutants including acid rain and the formation by free radical reaction of hazardous inorganic and organic compound e.g., PAN. (Analyzing)
- Describe causes and impacts of urban smog. (Analyzing)
- Explain greenhouse effect and global warming as resulting in climate change. (Analyzing)
- Explain the buildup to and recognize the adverse effects of ozone in the troposphere. (Applying)
- Describe the role of CFCs in destroying ozone in the stratosphere. (Applying)
- Describe the role of ozone in the stratosphere in reducing the intensity of harmful UV radiation reaching the earth. (Understanding)
- List possible alternatives to the use of CFCs. (Applying) Recognize and describe various water pollutants. (Applying)
- Explain the various parameters of water analysis. (Applying)
- List some major products of the petrochemicals together with their uses. (Applying)

It is the branch of chemistry which deals with the chemicals and other pollutants in the environment resulting directly and indirectly from human activities.

In this we study the sources, chemical reactions, transportation of the chemicals and their adverse effects on human beings.

Q1. What are different components of environment?

Answer

Components of the Environment:

- 1) Atmosphere → We will study it in detail
- 2) Hydrosphere → Concerned with all bodies i.e. ocean, rivers, streams lakes, glaciers and ground water reservoirs.
- 3) Lithosphere → Concerned with hard and rigid rocky earth crust.
- 4) Biosphere → Area on earth with support life i.e. air, lakes, etc.

Atmosphere: Our surrounding on earth is called atmosphere. It consists of gases i.e.

N_2 , CO , He, Ne, Kr, Xe and water vapours. Its thickness is about 1000 km above the surface of earth.

The gases present in the atmosphere are very important in the following ways:

- 1) These gases absorb harmful radiations (cosmic rays and electromagnetic radiation) of Sun to protect life on earth. Otherwise these rays are very harmful to living things on earth.
- 2) N_2 is used by nitrogen fixing bacteria
- 3) O_2 is necessary for breathing in animals.
- 4) CO_2 is necessary for photosynthesis in plants and
- 5) Water vapours are responsible for sustaining life on earth.
- 6) Actually our atmosphere has been divided into four layers:
 - 1) Troposphere
 - 2) Stratosphere
 - 3) Mesosphere
 - 4) Thermosphere.

We are concerned here with only first two layer i.e. troposphere and stratosphere.

Q2. Write a note on chemistry of troposphere.

Answer

Troposphere is very close to earth in which we live. It extends up to 20km. it contains all those gases which are present in our atmosphere.

In this part of our atmosphere, we will discuss

- 1) Different pollutants (Their sources and effects and smog (development and chemistry)
- 2) Effects and chemical reactions occur during acid rain, green effect and global warming
- 3) Role of automobile in air pollution

Smog

"It is a combination of smoke and fog i.e. 'sm' from smoke and 'og' from fog"

Under the right conditions, the smoke and sulfur dioxide produced from the burning of coal can combine with fog to create industrial smog.

Photochemical smog is a condition that develops when primary pollutants (oxides of nitrogen and volatile organic compounds created from fossil fuel combustion) interact under the influence of sunlight to produce a mixture of hundreds of different and hazardous chemicals known as secondary pollutants. Major Chemical Pollutants in Photochemical Smog: (Sources and Environmental Effects).

Toxic Chemical	Sources (Natural and Human)	Environmental Effects	Additional Notes
Nitrogen Oxides (NO and NO₂)	<ul style="list-style-type: none"> - Combustion of oil, coal, gas in both automobiles and industry. - bacterial action in soil - forest fires - volcanic action - lightning 	<ul style="list-style-type: none"> - Decreased visibility due to yellowish color of NO₂ - NO₂ contributes to heart and lung problems - NO₂ can suppress plant growth - decreased resistance to infection - may encourage the spread of cancer 	<ul style="list-style-type: none"> - All combustion processes account for only 5 % of NO₂ in the atmosphere, most is formed from reactions involving NO - concentrations likely to rise in the future
Volatile Organic Compounds (VOCs)	<ul style="list-style-type: none"> - evaporation of solvents - evaporation of fuels - incomplete combustion of fossil fuels - naturally occurring compounds like terpenes from trees 	<ul style="list-style-type: none"> - eye irritation - respiratory irritation - some are carcinogenic - decreased visibility due to blue-brown haze 	<ul style="list-style-type: none"> - the effects of VOCs are dependent on the type of chemical - samples show over 600 different VOCs in atmosphere - concentrations likely to continue to rise in future
Ozone (O₃)	<ul style="list-style-type: none"> - formed from photolysis of NO₂ - sometimes results from stratospheric ozone intrusions 	<ul style="list-style-type: none"> - bronchial constriction - coughing, Sneezing - respiratory irritation - eye irritation - decreased crop yields - retards plant growth - damages plastics - breaks down rubber - harsh odor 	<ul style="list-style-type: none"> - concentrations of 0.1 parts per million can reduce photosynthesis by 50 % - people with asthma and respiratory problems are influenced the most - can only be formed during daylight hours
Peroxyacetyl Nitrates (PAN)	<ul style="list-style-type: none"> - formed by the reaction of NO₂ with VOCs (can be formed naturally in some environments) 	<ul style="list-style-type: none"> - eye irritation - high toxicity to plants - respiratory irritation - damaging to proteins 	<ul style="list-style-type: none"> - was not detected until recognized in smog - higher toxicity to plants than ozone
	<ul style="list-style-type: none"> - Incomplete combustion of 	<ul style="list-style-type: none"> - Causes Anoxia (Oxygen 	

Carbon Oxide (CO and CO₂)	carbon containing fuels. (CO) <ul style="list-style-type: none"> - Incomplete combustion of agricultural of slush matter. (CO) - During the reaction in blast furnace (CO) - Cigarette smoke (CO) - Burning of fossil fuels produce CO₂ - Cultivation of Soil (CO₂) - Eruption of Volcanoes (CO₂) - Respiration of living organism. (CO₂) 	Starvation) result suffocation <ul style="list-style-type: none"> - Causes Green house and Global Warming results climatic changes - Causes Acid rain - Causes Acid rain <ul style="list-style-type: none"> - Respiratory irritation 	
Oxides of Sulphur (SO₂, SO₃)	<ul style="list-style-type: none"> - Volcanic Eruption - Burnt of oil & coal. - Sulphide ores are roasted 	<ul style="list-style-type: none"> - Causes acid rain - Respiratory irritation. - Loss of green color in plants - Fading in color of Fabrics, leathers, paper and paints. 	

Q3. Give Chemistry of Photochemical Smog (Chemical Reactions in the Atmosphere)


Answer

The previous section suggested that the development of photochemical smog is primarily determined by an abundance of nitrogen oxides and volatile organic compounds in the atmosphere and the presence of particular environmental conditions. To begin the chemical process of photochemical smog development the following conditions must occur:

- 1) Sunlight.
- 2) The production of oxides of nitrogen (NO_x).
- 3) The production of volatile organic compounds (VOCs).
- 4) Temperatures greater than 18 degrees Celsius.

If the above criteria are met, several reactions will occur producing the toxic chemical constituents of photochemical smog. The following discussion outlines the processes required for the formation of two most dominant toxic components: ozone (O₃) and peroxyacetyl nitrate (PAN). Which is primarily created from volatile organic compounds.

Nitrogen dioxide can be formed by one of the following reactions. Notice that the nitrogen oxide (NO) acts to remove ozone (O₃) from the atmosphere and this mechanism occurs naturally in an unpolluted atmosphere.



Sunlight can break down nitrogen dioxide (NO₂) back into nitrogen oxide (NO).



The atomic oxygen (O) formed in the above reaction then reacts with one of the abundant oxygen molecules (which makes up 20.94 % of the atmosphere) producing ozone (O₃).



Nitrogen dioxide (NO₂) can also react with radicals produced from volatile organic compounds in a series of reactions to form toxic products such as peroxyacetyl nitrates (PAN).



It should be noted that ozone can be produced naturally in an unpolluted atmosphere. However, it is consumed by nitrogen oxide as illustrated in the first

reaction. The introduction of volatile organic compounds results in an alternative pathway for the nitrogen oxide, still forming nitrogen dioxide but not consuming the ozone, and therefore ozone concentrations can be elevated to toxic levels.


Q4. What is Acid Rain? Give its harms.

Answer

Air contains SO_2 , NO_2 and CO_2 .

1) SO_2 present in air undergoes photolytic and catalytic oxidation to form SO_3 which reacts with rainy water or moisture to form H_2SO_4 i.e. $\text{SO}_3 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{SO}_4$

2) NO_2 reacts with rainy water or moisture in the presence of O_2 and O_3 and produces



3) CO_2 reacts with rainy water or moisture to form H_2CO_3 i.e. $\text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{CO}_3$

4) In some countries due to release of HCl by volcanic eruption, there is temporary acid rain.

H_2SO_4 , HNO_3 and H_2CO_3 formed as above come down the atmosphere as acid rain or acid snow. As far as H_2CO_3 is concerned, it has no severe effects on animals, plants or any other things.

Impacts:

1) It makes the lakes so acidic that they can no longer support fish life.

2) The yield of agricultural crops is also reduced.

3) HNO_3 acid rain gradually eats up lime stone and marble of the buildings and corrodes metals.

4) It fades the color of fabrics (e.g. cotton, nylon and rayon), leather and paper

5) Causes extensive leaf-drop in plants.

6) It is very corrosive and attacks skin.

7) Acidification of soil and rocks can leach metals like Al, Hg, Pb and Ca and discharge them into water bodies. Then these heavy metals are eaten by fishes which proves very much dangerous for those animals and birds which eat these fishes.

8) It also damages steel, paint, plastic, cement, masonry work and sculptural materials.

Q5. What is Green House Effect and Global Warming?

Answer

There is a protective layer O_3 gas in the atmosphere at a height between 15 km and 60 km. (The thickest layer of O_3 exists at a height of 23 km from the surface of the earth) and then a blanket of CO_2 gas exists in lower part of the atmosphere (i.e. below 15 km). Now when sunlight, consisting of ultra-violet rays, visible light and infra-red rays falls on the top of the atmosphere, the harmful ultra-violet rays are absorbed by the O_3 layer and hence do not reach the earth's surface. On the other hand, the visible light and infra-red rays pass through the CO_2 layer and fall on the earth. Since the infra-red radiations have heating effect, they heat the earth and its various objects.

Now since the earth and its various objects become hot (of course, less hot than the sun), they start emitting infra-red rays (heat rays), which are of long wavelength (note that the infra-red ray emitted by the extreme hot bodies like sun are of short wavelength). The infra-red radiations of long wavelength emitted by the earth and its objects are absorbed by the CO_2 layer in the atmosphere. Thus, we see that the presence of CO_2 in the atmosphere does not allow the infra-red radiation reflected by the earth's surface to go out of the atmosphere. In other words, we can say that the layer of CO_2 gas in the atmosphere traps all the infra-

red radiations coming from the earth's surface. These trapped infra-red rays heat the earth's atmosphere. The heating up of earth due to the trapping of infrared radiations (reflected from the earth's surface by CO_2 layer in the atmosphere is called "green-house-effect". Since the infrared rays coming from the earth cannot pass through the CO_2 layer, the temperature of the earth is raised. The rise in the temperature produced by the green-house-effect in the earth's atmosphere depends on the amount of CO_2 gas present in the atmosphere. The name green-house effect comes from the fact that this effect is used in horticulture for the upbringing of green plants in a small house, whose walls and roof are made of glass-sheet. The glass walls and glass roof of the house allow the short wavelength infrared radiations, contained in sunlight, to go into green-house freely, but do not allow the long wavelength infrared radiations, reflected by the soil, plants and other contents of the green-house to go out. These trapped infrared radiations raise the temperature inside the green house. Thus, even without the internal supply of heat, the temperature inside the green house higher than that outside it.

Importance of green-house effect:

The green house effect, produced by the presence of CO_2 layer in the atmosphere is, very necessary for our existence on the earth. We have seen that, due to green house effect CO_2 gas in the atmosphere does not allow the long wavelength infrared

radiations reflected by the earth, to go out of the atmosphere and hence the of the earth's and its atmosphere is increased. The rise in temperature of the earth is very necessary) for our existence on earth, because without it the whole earth would be converted into extremely cold planet and consequently we shall not be able to have a normal life.

Effect of excess of CO_2 present in the atmosphere:

If the atmosphere contains too much quantity of CO_2 , the green house effect is considerable increased i.e. due to excess quantity of CO_2 present in the atmosphere; the temperature of the earth is increased too much. This much high temperature melts all the glaciers (sno-mountains) floods the low-lying areas of the earth, changes the biological Activity) of oceans and the patterns of cropping etc. Thus we see that the presence of the excess of CO_2 in the atmosphere brings about climate changes.

Q6. Give pollutants produced by automobiles. What controls measures have made modern science in auto-mobiles?

Answer

In automobiles during incomplete combustion of petrol, we get smoke of different gases pollute atmosphere the engine used in these motor are called internal combustion) engines because the petrol which is used as a fuel is burnt inside the engine separately contains CO , NO , NO_2 , un-burnt carbon particles, some alcohol and acids. All these substances are poisonous and hence pollute air.

Production of Pollutants

Now let us see how they (CO , NO , NO_2) pollutants are produced in an engine of a car (automobile). Petrol is used as a fuel in automobile (car) engine. The main components of petrol are hydrocarbons. They hydrocarbons have general formula C_8H_{18} and hence are called octane.

Petrol burns very fast in a car engine. Due to short time available for burning, incomplete combustion of petrol takes and CO , unburnt particles, CO_2 , water some alcohol and acids are produced. CO and particles are emitted the air and thus air is polluted.

When petrol burns in a car engine, a very high temperature is produced. At this high temperature, N₂ and O₂ present in air of engine combine together and form NO and NO₂. The gases pollute the air.

Control Measure:

In order to control the air pollution caused by hydrocarbons and CO etc the following methods used.

1) By adding lead tetraethyl Pb(C₂H₅)₄, to petrol:

Suppose a used petrol as fuel. Petrol burns very fast in the vehicle engine, e.g. the rate of combustion of petrol is very high and hence the time taken by the petrol for its combustion is short. Due to this short time, the combustion of petrol is incomplete.


Pb(C₂H₅)₄ is added to the petrol to slow down the rate of combustion of provides Pb(C₂H₅)₄ more time to the petrol for its combustion and hence enables it to burn more completely'. Due to this the quantity of unburnt hydrocarbons and CO coming out of the engine, with exhaust gases is reduced and hence air pollution is also minimized. Pb is oxidized to PbO which gets deposited on the spark plugs and valves. In order to prevent the deposition of PbO, suitable amount of C₂H₄Cl₂ and C₂H₄Br₂ are also added to the petrol along with Pb(C₂H₅)₄. These halides convert PbO into PbCl₂ and PbBr₂ both of which are volatile and hence come out along with the exhaust gases emitted by the vehicle engine.



Although the addition of minimize the air pollution caused by unburnt hydrocarbons and CO, the air gets polluted with dangerous PbCl₂ and PbBr₂.

2) Catalytic oxidation/Converter:

The pollution of air caused by unburnt hydrocarbons and CO present in the exhaust gases of the vehicle's engine can also be reduced by attaching gas device with the vehicle's engine, in which the exhaust gases can be mixed with more air and then burnt completely in the presence of platinum catalyst before they are discharged into the environment. Hydrocarbons and CO are oxidized by O₂ of the air in presence of platinum catalyst.



The disadvantage of this method is that the platinum catalyst gets poisoned by PbCl₂ and PbBr₂ which are produced by the reaction between Pb(C₂H₅)₄, C₂H₄Cl₂ and C₂H₄Br₂ (these compounds are added to the petrol) in presence of CO₂.

Q7. Give chemistry of the stratosphere in detail.

Answer

This layer of atmosphere is present 20-40 km above the earth and ozone is present in this layer at a height of about 28 kilometers. The concentration of ozone in this layer is 10 ppm (10 parts per million)

In this layer of atmosphere, we will discuss comprehensively about ozone.

a) Production of O₃ and its toxic effects

Some O₃ is produced during various combustion processes taking place in the air around us. Traces of O₃ in air do not harm but O₃ of concentration more than 0.1 ppm is toxic and harmful to human beings. O₃ also attacks rubber products. O₃ is also produced in the upper part of the atmosphere by the action of sunlight on O₂.



b) Protective action of ozone layer in the atmosphere

The thickest layer of O₃ exists at a height of 23 km from the surface of the earth. Since O₃ present in this layer absorbs harmful ultraviolet radiations coming from the sun. If these radiations reach the earth, they will cause skin cancer and will destroy the organic molecules necessary for life.

Thus, we see that O₃ does not allow the ultra-violet radiations to reach the earth and we


are thus saved from the harmful effects caused by these radiations. If O₃ layer in the atmosphere disappears completely, then all the harmful ultra-violet radiations coming from the sun would reach the earth and would cause skin cancer in men and animals and will also damage the plants. All the life on earth would then gradually be destroyed.

c) Sources of destruction/destroying the ozone layer present in stratosphere.

In 1980 scientists showed that there is a hole in the O₃ layer. This hole was detected over the region of Antarctica. Due to the presence of O₃ layer, the ultraviolet rays coming from the sun can pass through the hole and thus can reach the earth's surface. The presence of O₃ in the atmosphere is due to the fact that the amount of O₃ present in stratosphere is getting reduced day by day and thus the ozone layer is becoming thinner and thinner. The depletion of the ozone layer is due to the following sources.

1) Oxides of Nitrogen:

The oxides of nitrogen present in the atmosphere decompose O₃ into O₂ and are themselves regenerated.



Thus we see that the presence of nitrogen oxides in the atmosphere destroys the ozone layer. These oxides destroy about 70% of O₃ found in the stratosphere.

Greater is the amount of the oxides in the atmosphere greater is the percentage of O_3 which is destroyed.

2) Nuclear Tests:

Nuclear tests being conducted in the world generate high temperature. At high temperature, atmospheric nitrogen is favorably oxidized to NO. NO thus formed destroys ozone layer, as shown above at (I)

3) Use of Chloro-Flouro Carbons:

Fluoro-chloro-carbons are the fluoro-chloro methanes like, Freon-I ($CFCl_3$) and Freon 12 (CF_2Cl_2). These are stable compounds. These are chemically inert and hence do not react with the substances. These are used as aerosol spray propellants, refrigerants, firefighting reagent and solvents for cleaning electronic components. When they enter stratosphere, they absorb ultraviolet solar radiations and get broken down into free atomic chlorine. This atomic chlorine decomposes O_3 into O_2 (NO also breaks O_3 into O_2).



d) How to protect the ozone layer:

Scientists are worried over the gradual destruction of ozone layer by the oxides of nitrogen and fluoro carbons. In order to save the destruction of O_3 layer by fluorochloro carbons, their use should be banned or some new types of substances should be discovered which may be used as aerosol spray propellants and should not react with O_3 layer, so that it may be saved.

Some Alternatives to Chloro-Flouro Carbons (CFCs):

The first CFC substitutes to be introduced were HCFCs such as CF_3CHCl_2 and CHF_2Cl compounds that have fewer chlorine atoms than other CFCs. HCFCs break down more readily in the atmosphere than CFCs and thus are less likely to

reach the stratosphere. However, because they can cause some O₃ destruction, they are scheduled to be phased out by 2030.

Much better substitutes for CFCs are hydrofluoro carbons which contain no chlorine. One of them CF₃CH₂F has been used successfully as a refrigerant and since 1994 has replaced Freon in nearly all car air conditioners. In electronic industries, soapy water followed by rinsing and air drying is now used instead of CFCs to clean micro circuits. Unfortunately, a serious problem is associated with the long term use of HFC. Like CFC and HCFCs they contribute to climate change. However researches are going on to find chemicals that are both effective refrigerant and environmental friendly.

Q8. What is water pollution and water treatment?

Answer

The contamination of water with the substances which have adverse effects on human beings, animal and plants is called water pollution

And the substances whose presence in the water makes it polluted are called pollutants.

Types of Water pollutants

These are of following types:

Suspended Solid and Sediments:

These are wastes which are not completely soluble but suspended in the water.

These wastes include:

- | | |
|--------------------------|----------------------|
| 1) Oil spillage | 2) Live-stock waste |
| 3) Industrial wastes and | 4) Leather tanneries |

1) Oil Spillage:

Petroleum is a complex mixture of many compounds which are mainly hydrocarbons. It is transported from one place to another through sea. We know that petroleum products are used as:

- 1) Fuel
- 2) Lubricants manufacture of petrochemicals
- 3) Plastics
- 4) Electrical appliances
- 5) Synthetic rubber
- 6) Detergents

In order to prepare such a large variety of substances, petroleum is handled on large scale in the world. So, the oil spillage can take place and it creates serious problems.

Pollution of water by petroleum:

Water gets polluted by:

- 1) accidental oil spills
- 2) leakage from cargo oil tankers in sea
- 3) tanker trucks
- 4) pipelines leakage during offshore exploration
- 5) leakage of underground storage tanks

Oil Spillage and animal life:

Many petroleum products are poisonous and create serious health problems to:

- 1) Humans
- 2) Animals
- 3) Aquatic life

Polycyclic hydrocarbons are carcinogenic even at very low concentration. Marine animals are seriously affected by soluble aromatic fractions of oil. The spilled oil damages the feather of the birds or fur of animals and sometimes causes their death.

Petroleum and under water plants:

When oil is spilled on the surface of sea then the light transmission is affected. The process of photosynthesis of plants does not remain much efficient moreover, the concentration of oxygen in water is decreased.

2) Live-Stock Waste:

Livestock waste is dumped on open land. Sometimes it is discharged into sewage, canals or rivers. This practice pollutes the surface and ground water. In this way serious problem are created for the population.

Bacteria are present in the livestock waste. It contaminates the surface and ground water. This causes the diseases like:

a) Dysentery

b) Typhoid

c) Hepatitis

3) Industrial Wastes:

Sources:

Industries which are producing large quantities of industrial effluents are leather, tanneries, fertilizers, oil refineries, petrochemicals, textiles, foods, sugar, paper, pulp, paper board, rubber products etc. the waste products may be waste heat, smoke solid or water effluent.

Pollutants:

The industrial pollutants are highly toxic organic compounds and heavy metals like Pb, Cd, Cr, Hg, As, Sb etc. Oil greases, Mineral acids are also release in small quantities. These pollutants result in contamination of water and make it unsuitable for irrigation and drinking purposes.

Effects of Industrial pollutants:

Heavy metal particles are highly toxic and do not have any safe limits. When they are continuously ingested through food or water they get accumulated in the organisms and cause serious health problems like anemia, kidney diseases, nervous disorder, high blood pressure etc.

4) Leather Tanneries:

There are many leather tanning units working in Pakistan. Their sizes vary from cottage scale to big industrial units.

Leather industries use chromium salts which have +6 oxidation state of chromium. Only a few industries have the facility of waste treatment. This can be done by reducing Cr⁺⁶ oxidation states to Cr⁺³ oxidation state. Cr⁺³ is precipitated as Cr(OH)₃. Cr⁺⁶ salts are highly toxic and cause cancer.

Dissolved Solids

These are wastes which are dissolved/soluble in water completely. These wastes include:

- 1) Detergents
- 2) Pesticides
- 3) Chemical fertilizers

1) Detergents:

Detergents are used in homes and industries for washing. After washing these detergents are thrown into water reservoirs. The amount of detergents in reservoirs is increasing day by day. This waste water containing detergents goes into rivers and finally reaches the ocean; which is harmful for life in seas. The detergents bound heavy metal ions like Pb, Cd and Hg and transport it from sediments into water. These metals particles are very toxic.

2) Pesticides:

The pesticides are both toxic and persistent. Analysis of polluted water has shown that it contains pesticides which are toxic to fish. Endrin, even in traces, is reported to be toxic for catfish and other varieties of fish. DDT, affects the central nervous system of fish and toxaphene has been reported to cause bone degeneration in fish.

3) Chemical Fertilizers:

Nitrate/phosphate salts are generally used as fertilizers, to increase the yield of the crops. When these fertilizers are used in excess, some of their unused quantity is washed away from the agriculture lands into the ponds, lakes and river with rain water and thus pollute the water. This water is polluted, since it contains unused nitrate/phosphate salts. Similarly, the waste water coming from the fertilizer industries also contains nitrogenous/phosphatic fertilizers which, when washed away into the lakes and river with rain water, make the water polluted. The presence of nitrogenous/phosphatic fertilizers in water is harmful to the aquatic life and human beings in the following ways:

- a) The presence of the fertilizers in the polluted water increases the growth of algae and other aquatic plants which, later on undergo decomposition and produce disagreeable odor. These plants also deplete the amount of O_2 dissolved in water and hence the survival of aquatic life becomes difficult or impossible.
- b) After a long period, the lakes and slow-moving waters which contains plant nutrients are converted into swamps (A swamp is an area of very wet land with wild plants growing in it) and marshes (A marsh is an area of land which is very wet and muddy).
- c) The water containing nitrate salt is not fit for drinking by human beings. Moreover, this polluted water cannot be purified for drinking purposes.

Thermal Pollution

Thermal pollution takes place because many electric generating companies use water in the process of cooling their generator. This heated water is then


released into the system from which it was drawn, causing a warming trend of the surface, water. Thermal pollution results when the heated effluent is released into poorly flushed system. In these cases, permanent temperature increase often result, which tend to decrease the solubility of dissolved oxygen. In lakes it also becomes possible to bring about nutrient redistributions and prolong summer stagnation periods.

When heated water gets released into large, well-flushed marine systems there is little if any permanent temperature rise. There are however problems related to the operation of plants utilizing marine water in the cooling process. Evidence reveals that seawater tends to corrode the cooling pipes, which are generally constructed of a copper nickel alloy termed Monel. These metals readily dissolved in the heated seawater and are then released into the marine environment together with the heated effluent. This adds to the nickel and copper concentrations of these systems. In addition, the screens covering the water intake pipes rapidly foul with marine organism, which decrease the flow of water into the plant. The screens have been commonly cleaned by using a concentrated detergent solution or copper sulphate. These cleaning materials have been then released into the contaminated the surrounding waters.

Q9. Write a detailed note on: Waste Water Treatment

Answer

i) Analysis/Purification of Water



Industrial wastewater treatment

Covers the mechanisms and processes used to treat waters that have been contaminated in some way by man's industrial or commercial activities prior to its release into the environment or its re-use.

Most industries produce some wet waste although recent trends in the developed world have been to minimize such production or recycle such waste within the production process. However, many industries remain dependent on processes that produce water based waste stream.

Treatment of industrial wastewater

The different types of contamination of wastewater require a variety of strategies to remove the contamination.

a) Solids removal

Most solids can be removed using simple sedimentation techniques with the solids recovered as slurry or sludge.

b) Oils and greases

Many oils can be recovered from open water surfaces by skimming devices. However, hydraulic oils and the majority of oils that have degraded to any extent will also have a soluble or emulsified component that will require further treatment to eliminate.

c) Soft organics

Organic material of plant or animal origin is usually possible to treat using extended conventional Wastewater treatment processes. Problems can arise if the wastewater is excessively diluted with washing water or is highly concentrated such as neat blood or milk. The presence of cleaning agents, disinfectants, pesticides, or antibiotics can have detrimental impacts on processes.

d) Hard organics

Synthetic organic materials including solvents, paints, pharmaceuticals, pesticides, coking products etc can be very difficult to treat. Treatment methods are often specific to the material being treated. Methods include distillation, adsorption, nitrification, incineration, chemical immobilization or landfill disposal.

e) Acids and alkalis

Acids and alkalis usually be neutralized under controlled conditions. Neutralization frequently a precipitate that will require treatment as a solid residue that may also be toxic. In some cases, gasses may be evolved requiring treatment for the gas

f) Toxic materials

Toxic materials including many organic materials, metals (such as zinc, silver, cadmium, thallium etc.) acids, alkalis, non-metallic elements (such as arsenic or selenium) are generally resistant to biological processes unless very dilute. Metals can often be precipitated out by changing the pH or by treatment with other

chemicals. Many, however, are resistant to treatment or mitigation and may require concentration followed by land filling or recycling.

Various Parameters of Water Analysis

Following table will help us to determine the water quality.

Table Water — Quality Indicators

Parameter	Significance	Level
Dissolved oxygen	General indicator of water quality; source of O ₂ for respiration	Minimum acceptable level, 4-5 mg/liter; 10-15 mg/liter for reproduction of desirable fish
Total suspended solids	Clog fish gills, bury eggs reduce light penetration increase heat absorption	Dependent on location
Total dissolved solids	Represents total mineral content which may or may not be toxic	A maximum of 400 mg liter for diverse fish populations
BOD	Amount of dissolved oxygen removed during decomposition of organic matter in a given time; a general indicator of contamination due to biodegradable organics	BOD Water Status 1 mg/liter Very clean 2 mg/liter Clean 5 mg/liter Fairly clean 10 mg/liter Doubtful Contaminated
COD	Indicates the concentration of materials oxidizable by chemical reaction	0.5 mg/liter indicates very clean streams

pH	Indicates the addition of acid or bases	pH depends on actual system
Iron	Excessive amounts can clog fish gills; indicates drainage from iron bearing sediments, mines and industrial processes	A maximum of 0.7 mg/liter for diverse fish populations
Manganese	Concentration low in natural systems due to low solubility high concentrations indicates contamination	maximum of 1 mg/liter is common criterion for quality

Table (Contd.)

Parameter	Significance	Level
Copper	Indicates drainage from copper-bearing sediment, mines, plating, or other industrial sources	A maximum of 0.02- 10 mg/liter is a common criterion for stream quality
Zinc	Indicates mine drainage or industrial input	A maximum of 1 mg/liter is common criterion for stream quality
Hg, Cd, Pb, Ni, Cr, Ag, etc.	Indicates industrial input	A maximum of 1 mg/liter is common criterion for stream quality
Nitrate	A major plant nutrient; in high-concentration it can promote excessive plant growth; major sources are fertilizers, sludge, and sewage	A maximum of 0.3 mg/liter to prevent excessive fertilization of streams

Phosphate	A major plant nutrient; major sources are detergents, fertilizer, sewage	A maximum of 0.03-0.4 mg/liter total inorganic phosphate is a common criterion
-----------	--	--

Q10.What is Green Chemistry? Give its importance.**Answer**

The term green chemistry, coined in 1991, is defined as "the design of chemical products and processes that reduce or eliminate the use and generation of hazardous substances."

Green chemistry emphasizes the design and creation of chemicals that are not hazardous to people or the environment. It has been applied to a wide range of industrial and consumer goods, including paints, dyes, fertilizers, pesticides, plastics, medicines, electronics, dry cleaning, energy generation, and water purification.

Why Green Chemistry?

Green chemistry is effective in reducing the impact of chemicals on human health and the environment. In addition, many companies have found that it can be cheaper and even profitable to meet environmental goals. Profits derive from higher efficiency, less waste, better product quality, and reduced liability. Many environmental laws and regulations target hazardous chemicals, and following all these requirements can be complicated. But green chemistry allows companies to comply with the law in much simpler and cheaper ways. Finally, green chemistry is a fundamental science-based approach. Addressing the problem of hazard at the molecular level, it can be applied to all kinds of environmental issues. •

Since 1991, there have been many advances in green chemistry, in both academic research and industrial implementation. For example, Spinosad, an insecticide manufactured by fermenting a naturally occurring soil organism, was registered by the EPA as a reduced-risk insecticide in 1997. Spinosad does not leach, bioaccumulate, volatilize, or persist in the environment and in field tests left 70 to 90 percent of beneficial insects unharmed. It has a relatively low toxicity to mammals and birds and is slightly to moderately toxic to aquatic organisms, but is toxic to bees until it dries.

The 12 Principles of Green Chemistry

The aim of green chemistry is to reduce chemical related impact on human health and virtually eliminate contamination of the environment through dedicated, sustainable prevention programs. Green chemistry searches for alternative, environmentally friendly reaction media and at the same time strives to increase reaction rates and lower reaction temperatures. The green chemistry concept applies innovative scientific solutions to solve environmental issues posed in the laboratory. Paul T. Anastas, an organic chemist working in the Office of Pollution Prevention and Toxins at the EPA, and John C. Warner developed the Twelve Principles of Green Chemistry in 1991. These principles can be grouped into "Reducing Risk" and "Minimizing the

Environmental Footprint."

I. Reducing Risk in the Laboratory

Sigma-Aldrich is dedicated to providing alternative products designed with the health and safety of its employees, customers, and the public in mind.

- i) Use Safer Chemicals — Utilize performance chemicals that have the lowest levels of toxicity.
- ii) Design Less Hazardous Synthesis Methods — Where feasible, make use of synthetic or biosynthetic methods that pose little or no toxicity to human health and the environment.

- iii) Use Safer Solvents and Reaction Conditions Search for the most up-to-date information on green solvents that will optimize your process and provide a safer working environment.
- iv) Accident Prevention — Select substances that minimize the potential for explosions, fires and chemical releases into the environment.

II. Minimizing the Environmental Footprint

The 12 Principles focus on reducing the volumes of chemicals used and pollution prevention.

- i) Waste Minimization and Prevention — Develop chemical synthesis techniques, which reduce or prevent waste. It is better to prevent waste than to clean it up after its creation.
- ii) Use of Catalysts Instead of Stoichiometric Quantities — Catalytic reactions inherently use smaller quantities of chemicals to carry out a specified transformation.
- iii) Reduce the Use of Chemical Derivatives — The use of protecting groups or other forms of temporary modification of a functionality adds to the total waste incurred in a synthetic route.
- iv) Synthetic Efficiency (Atom Economy) — An efficient chemical process ensures the maximum amount of your starting materials is used in the final product so that no atom is wasted.
- v) Taking Advantage of Chemicals Designed for Degradation — Reduce the effect on the environment by using chemicals that are designed to be biodegradable.
- vi) Establishment of In Process Controls for Pollution Prevention — To avoid the formation of hazardous substances, adopt real-time analysis and in process monitoring during synthesis.

vii) Use of Renewable Feedstocks — Use raw materials or renewable feedstocks (waste from other processes or products derived from agricultural streams) whenever technically or economically feasible.

viii) Encourage Energy Efficiency — The realization of the economic and environmental impact of energy use in a chemical process and the development of alternative means to reduce the impact.

Q11. What is radioactive radiation pollution? Give its control.

Answer

Radiation Pollution

What is radioactive pollution?

Radioactive substances and nuclear radiations produced during nuclear reaction affect our environment adversely and thus radioactive pollution created.

Sources of radioactive pollution and its effects

1. Low level radioactive liquid wastes, radioactive gaseous wastes, and dusts are released during nuclear explosions. The radioactive gaseous wastes are injected into the upper layer of atmosphere where, due to cooling they condense to fine dust particles and thus radioactive cloud is formed. This cloud moves in the direction of the wind, settles down slowly to the surface of the earth and thus pollutes air, water and soil.
2. The radioactive substances produce energy which is so strong that the living cells are damaged or destroyed.
3. People working with radioactive elements develop tumors.
4. Radioactive element like strontium 90 affects our soil and through this human beings and animals are also affected adversely.

5. Nuclear explosions which are operated in sea make sea water polluted. This affects the aquatic life.
6. Among the radioactive radiations, gamma radiations are the most dangerous, since they have high energy and big penetrating power. These radiations can, therefore, pass freely in the human body, where they lose energy, which destroys the living cells by converting them into charged particles (ions). These charged particles are chemically very reactive and hence disrupt cell membrane, reduce the effectiveness of enzymes and even damage genes and chromosomes. All this results in diseases like leukemia and cancer.
7. We know that in a nuclear reactor U 235 is used as a nuclear fuel, which undergoes nuclear fission and energy is produced. Nuclear radiations are produced in the processes viz mining and enrichment of U 235 taking place in the nuclear reactor. These radiations can leak from the reactor and therefore, damage the health of the human beings and animals.

Control measures for minimizing radioactive pollution

The waste material produced in the mining, enrichment and fission of U 235 inside the reactor are collectively called nuclear wastes. At present most of the nuclear wastes are being stored in strong leak proof containers. These will be disposed off whenever a safe method of their disposal is found out.

Using catalytic Converters

A catalytic converter removes pollutant gases from the exhaust by oxidizing or reducing them. The exhaust gases pass through a converter containing a precious metal catalyst, usually an alloy of platinum and rhodium. Several reactions may take place. NOX and CO may take part in a redox reaction which neatly removes both of them at the same time: NOX oxidizes CO to CO₂ and is reduced to harmless nitrogen gas.




CO and C_xH_x are oxidized by air;



(Using C_7H_{16} to represent a typical hydrocarbon.)

For all three of these reactions happen, it is necessary to use a three way converter and to have an oxygen monitor fitted to the engine, this checks the quantity of oxygen going into the engine to make sure there is enough to carry out the oxidation reactions.



The overall result of passing exhaust gases through this kind of catalyst system is to convert CO, NO_x and C_xH_y to relatively harmless N₂, CO₂ and H₂O.

The catalytic reactions do not start working until the catalyst has reached a temperature of about 200°C so they are not effective until the engine has warmed up.

Catalyst systems of this type cost several hundred pounds, mainly because of the high cost of the precious metal they contain. The catalyst is poisoned by lead, so unleaded fuel must always be used.

Q12. Why waste water from household and industries without treatment to river is dangerous?

Answer

When we think about river pollution, we might assume it comes from places like factories, farms and industry. Waste water from manufacturing or chemical processes in industries contributes to water pollution. Industrial waste water usually contains specific and readily identifiable chemical compounds yet, in many cases the pollution in our rivers comes from a much less obvious source - our homes. Incorrect plumbing could mean that waste water from dishwashers, washing machines, sinks, baths and even toilets other impurities includes organic materials and plant nutrients that tend to rot. The main organic materials are food and vegetable waste, plant nutrient come from chemical soaps, washing powders, etc. These flushed directly into a local river. These 'misconnected' pipes are a common cause of pollution to rivers and streams, especially in towns and cities.

There are normally two forms of drainage -surface water and foul water.

Surface water drains, or 'storm drains' carry rainwater from road surfaces and rooftops into local rivers and streams and flows into the river untreated.

Foul water drains carry waste water from toilets, sinks, baths and household appliances to the sewage treatment works. This water is treated before it can safely flow back into river and streams.

Today, many people dump their garbage into streams, lakes, rivers, and seas, thus making water bodies the final resting place of cans, bottles, plastics, and other household products. The various substances that we use for keeping our houses clean add to water pollution as they contain harmful chemicals.

Americans generate 1.6 million tons of household hazardous waste per year. The average home can accumulate as much as 100 pounds of household hazardous waste in the basement or garage and in storage closets. When improperly disposed of, household hazardous waste can create a risk to people and the environment. Paints, cleaners, oils, batteries, and pesticides are

examples of just a few of the common household hazardous wastes that need special disposal.

When fresh water is artificially supplemented with nutrients, it results in an abnormal increase in the growth of water plants. This is known as eutrophication. The discharge of waste from industries, agriculture, and urban communities into water bodies generally stretches the biological capacities of aquatic systems. Chemical run-off from fields also adds nutrients to water. Excess nutrients cause the water body to become choked with organic substances and organisms. When organic matter exceeds the capacity of the micro-organisms in water that break down and recycle the organic matter, it encourages rapid growth, or blooms, of algae. When they die, the remains of the algae add to the organic wastes already in the water; eventually, the water becomes deficient in oxygen. Anaerobic organisms (those that do not require oxygen to live) then attack the organic wastes, releasing gases such as methane and hydrogen sulphide, which are harmful to the oxygen-requiring (aerobic) forms of life. The result is a foul smelling, waste-filled body of water. Untreated sewage effluent in the water causes oxygen levels to drop drastically, sewage fungus covers the bed of the watercourse like a blanket and in more severe cases the river can no longer support fish, insects and animals that live in and around the water.

Polluted water is unsuitable for drinking, recreation, agriculture, and industry. It diminishes the aesthetic quality of lakes and rivers. More seriously, contaminated water destroys aquatic life and reduces its reproductive ability. Eventually, it is a hazard to human health. Nobody can escape the effects of water pollution.

Once an aquifer is contaminated, it may be unusable for decades. The residence time, as noted earlier, can be anywhere from two weeks or 10 000 years.

Q13. What are different alternatives to ozone-depleting halocarbons?**Answer**

Hydrofluorocarbons (HFCs) and perfluorocarbons (PFCs) are alternative halocarbons being used to replace ODS in industrial and commercial applications and their use is becoming more widespread. Other alternatives to ODS include ammonia, carbon dioxide and hydrocarbons.

The relationship between ozone-depleting substances and their halocarbon alternatives and global warming

Ozone-depleting substances (ODS) and their halocarbon alternatives have a direct global warming potential and contribute to climate change. The production of some ozone depleting substances (e.g. CFCs) has been ceased worldwide resulting in emission reductions of ozone-depleting substances into the atmosphere. However, emissions of other ODS (e.g. hydrochlorofluorocarbons [HCFCs]) and ODS alternatives (e.g. hydrofluorocarbons [HFCs]) is expected to continue to increase.

HFCs are ODS refrigerants alternatives and their use is becoming more widespread. Although HFCs have no ozone-depletion potential they do have high global-warming potential and are thousands of times more potent greenhouse gases than carbon dioxide.

Regulations that prevent or minimize ODS and other halocarbons emissions serve a dual environmental benefit of lowering emissions that destroy the ozone layer and contribute to climate change.

Q14. Are ozone-depleting substances still an important environmental concern?**Answer**

Although the production and use of many ODS has been phased out, the control of ODS emissions continues to be an important environmental issue, as the destruction of the earth's protective stratospheric ozone layer causes increased health risks and environmental impacts, such as cataracts, skin cancer and climate change.

Air Pollution

1. In past there was bull cart, donkey carts animals were used by but now a days Aeroplane, cars, buses and trains are used for transportations. These vehicles need energy for their working in the form of petrol or gas to provide energy for motion of vehicle. The end product of this burning includes energy plus different gasses include hydrocarbons, Nitrogen oxides carbon monoxide, Benzene, sulphur heavy metals and particles, these gases enter into air and pollute in when human breath in such polluted environment it causes asthma, coughing, nausea, chest pain, bronchitis diseases etc.
2. It is estimated that usually 24000 deaths occur due to air pollution because pollution damage central Nervous System.
3. Automobiles are responsible for depletion of ozone in stratosphere sheet because of source of CFCs entering the atmosphere which is escape from vehicle air condition (ACS)

Vehicles are responsible for 80% pollution in metropolitan region three fold increases in per capita energy consumption for global warming as burning of fuel produces CO_2 .

4. Ways to improve impacts of car like uses of small cars, cleaner cars, efficient engine catalytic converters are compulsory and use of biofuel.
 - A. Fuel cell causes only water comes out as its exhaust electrical vehicle pose problem because of range of batteries and sepe. Hybrid electric petrol they are providing solution for this pollution they cut down the fuel wastage and effect the high energy fuel on board e.g. in Delhi they demolished the buses run on diesel

B. Restricts the occurs to the cities or part of cities e.g. Chester York, oxford have driven the apparently to park ad ride they have given free space for parking the edge of town.

C. improve information about road condition and public transport e.g. Zwich zone referendum eh priority is given to tarns trolleys, buses, parking prohibitions 50% of all the trips as by means of public transport

Polluter pay is another solution where the pollution emitted by the car is measured and the driver is charged on the buses of level multiplies the distance travel.

Technological changes to reduce cooling zone of engine wall and reduce hydrocarbon and carbon monoxide emission. Computer control at IC engine.

Catalytic converter technology (CCT) emission

Trap oxidizers ad ceramic filter a Diesel vehicles

Direct injection fuel efficient Diesel

Ozone

Ozone layer is found in the stratosphere approximately 10-50 Km above the earth surface. Ozone molecules have three atoms of oxygen instead of the normal tow. The ozone layer protects us from the harmful effects of certain wave lengths of ultra violet (UV) light coming from the sun, socially UVB. Any significant decrease in ozone in the atmosphere would result in an increase of UV.B radiation reaching the earth surface. Increase in levels at UV.B radiation can result in the increase in skin cancer; suppress the immune system, eye disorder including catavart and effect plants, animals and plastic materials.

Ozone Depletion

In 1985, scientists discovered that there is severe ozone depletion in the Antarctic region, which was confirmed by American Satellite observations.

Chlorofluorocarbons (CFCs) were invented in 1928 found many used in foams, refrigeration, air conditioners, solvent, fire extinguishers etc. these CFCs are long lived and their emissions reach stratosphere and cause ozone depletion. This has been dramatically confirmed through Antarctic ozone hole.

The United Nations Environment Programme (UNEP) has been addressing the issue of depletion of the ozone layer since 1977 and in 1981 UNEP's Governing Council set up a working group to prepare nine global framework conventions for protection of ozone layer.

By reducing the use of CFCs we can protect ozone layer for saving the environment from harmful effects.

Montreal protocol on substances that deplete the ozone layer was finally agreed upon on 16th September 1987 and adopted by the Government in this protocol ozone depleting substances are banned.

Q15. What are different natural water purification methods?

Answer

There are several, natural ways that you can purify water.

In the event of an emergency it may become necessary to purify the water that you drink to avoid getting sick. Though many people think purifying water is difficult, there are a number of simple, natural means of water that can easily be used in an emergency.



1) Sand

One of the most common ways that water is purified is by passing it through sand and soil. If contaminated water (say urine) is poured onto fine sand, the

water is going to travel down until it reaches the saturation level. The other chemicals that are mixed with the water, though, will be held back by the sand until only the water is left. This is the same method of water purification used in many sewage treatment facilities as a big part of separating water out of the waste that goes through the plant.

2) Boiling

Water can be boiled to make it purer. Boiling water kills bacteria and other germs in the water, which makes it safer to drink. Or, build a water still that functions off boiling. Water is poured into a sealed pot, with a tube leading out of it. The water turns into steam, and the steam escapes through the tube. The tube transports the steam to another container, where the steam condenses back into water. This process can eliminate more contaminants from water (such as salt), but it's also more complicated.

3) Cloth Filtration

Cloth filters can be used to help keep larger contaminants out of water. This process is very simple; pour the water through a cloth, or through several layers of cloth, and the cloths' weave will strain the water and hold back impurities. This is the same process that happens when a coffee filter is used, except that the filter is paper rather than cloth. Paper can work as well, and if you have coffee filters or similar devices you can easily use those instead of a clean cloth to strain your water.

Q16. How rain water seepage through hazardous waste dumpsites can dissolve in drinking water supplies Hazardous waste:

Answer

Hazardous waste is waste that is dangerous or potentially harmful to our health or the environment. Hazardous wastes can be liquids, solids, gases, or sludges.

They can be discarded commercial products, like cleaning fluids or pesticides, or the by-products of manufacturing processes. It's very important to dispose off hazardous waste carefully otherwise it contaminates the air, water, and soil.

Groundwater:

Groundwater is rain water or water from surface water bodies, like lakes or streams, that soaks into the soil and bedrock and is stored underground in the tiny spaces between rocks and particles of soil.

Groundwater contaminants come from two categories of sources: point sources and distributed, or non-point sources. Landfills, leaking gasoline storage tanks, leaking septic tanks, and accidental spills are examples of point sources.

Infiltration from farm land treated with pesticides and fertilizers is an example of a non-point source.

1. Among the more significant point sources are municipal landfills and industrial waste disposal sites. When either of these occur in or near sand and gravel aquifers, the potential for widespread contamination is the greatest.
2. Leaks of petroleum products have been increasing over the last two decades because underground steel tanks installed in large numbers in the 1950s and 1960s have become corroded. Before 1980, most underground tanks were made of steel. Without adequate corrosion protection, up to half of them leak by the time they are 15 years old. Groundwater dissolves many different compounds, and most of these substances have the potential to contaminate large quantities of water. For example, one litre of gasoline can contaminate 1 000 000 litres of groundwater. This problem is particularly severe in the Atlantic provinces where there is a high usage of groundwater. In many cases, the problem is noticed long after the aquifer is contaminated, for example, when consumers start tasting or smelling gasoline.

3. Groundwater can become contaminated in many ways. Chemicals from hazardous wastes buried in unsecured landfills. If rain water or surface water comes into contact with contaminated soil while seeping into the ground, it can become polluted and can carry the pollution from the soil to the groundwater. From here, contaminants can spread to wells or surface water, making it unsafe to drink.
4. Groundwater can also become contaminated when liquid hazardous substances themselves soak down through the soil or rock into the groundwater. Some liquid hazardous substances do not mix with the groundwater but remain pooled within the soil or bedrock. These pooled substances can act as long-term sources of groundwater contamination as the groundwater flows through the soil or rock and comes into contact with them. Groundwater contamination is extremely difficult, and sometimes impossible, to clean up.
5. Apart from chemical pollutants the major culprits are Bacteria and Viruses which cause most of the commonly found water borne diseases.
6. Bacterial diseases: Gastro-enteritis, Typhoid, Cholera, Paratyphoid, Dysentery and Diarrhea.

Viral diseases: Polio, Dysentery, Gastro-enteritis, Diarrhea and Jaundice (Hepatitis)

Replacing Chlorofluoro Carbons

Few compounds have CFCs combination of non-flammability, non-toxicity and inertness, and for uses such as refrigeration and aerosols it is necessary to find compounds with exactly the right boiling point.

Some of the important replacements for these uses are the hydrofluorocarbons, CFCs (also known as hydrofluorocarbon or HFAs.). An example is 1,1,1,2-tetrafluoroethane, $\text{CH}_3\text{CF}_2\text{F}$, which is used as a refrigerant. HFCs have the advantage that they contain no Cl atoms, so they do not release damaging Cl radicals in the stratosphere. Moreover, their molecules include C-H bonds, which

are relatively reactive, which means that these compounds break down in the atmosphere more quickly than CFCs so they do not persist for so long.

